

CHRONIC INFLAMMATION AFTER CATARACT SURGERY



Tips on the management and comanagement of these cases from a retina specialist.

BY VAIDEHI S. DEDANIA, MD

Chronic or recurrent inflammation, including cystoid macular edema (CME), occurs in about 0.1% to 2% of patients following routine cataract surgery.¹ Prolonged postsurgical inflammation, though relatively rare, can be frustrating for both patients and practitioners. Risk factors include a history of uveitis or diabetes and long and/or complicated surgery.² In some cases, an indolent infectious organism introduced at the time of surgery leads to chronic or recurrent postoperative inflammation.³

Practicing at an academic referral center, I regularly comanage patients who develop chronic inflammation after cataract surgery. Most of these individuals present to my practice 6 to 12 weeks postoperatively—sometimes because the referring surgeon detected CME with OCT and sometimes because the patient reported blurred vision, photophobia, and/or pain. This article shares my tips on successful management and comanagement.

DIAGNOSIS

I choose diagnostic tests depending on the patient's status upon referral. If the individual has relatively mild inflammation and a straightforward history, then topical steroids are restarted and tapered over a 3-month period. If the inflammation is severe, then a workup is performed to rule out other infectious and inflammatory etiologies, particularly before an intraocular steroid injection is considered. The examination includes a uveitis workup and, on a

case-by-case basis, an anterior chamber tap for polymerase chain reaction or culture. A workup is also in order if something such as plaque is observed on the lens that may be suspicious for infection. Much of the time, the inflammation presents primarily in the anterior chamber; I have a lower threshold for a complete uveitis workup if vitreous haze is evident.

Some patients who experience chronic or recurrent postoperative inflammation after cataract surgery on their first eye develop it after cataract surgery on their second eye. Generally, if any of the aforementioned features are present, then I perform an additional evaluation. Otherwise, I bear in mind that these patients may have a predisposition to inflammation due to unidentifiable factors.

TREATMENT ALGORITHM

If a patient is diagnosed with chronic or recurrent postoperative inflammation 6 to 8 weeks after cataract surgery, then the postoperative steroid is typically restarted (usually at a higher dose). If the inflammation is severe, then dosing may be as frequent as every 2 hours for a few days before being tapered. If the inflammation is mild to moderate, then the steroid is administered four times per day. The presence of CME may warrant an extended taper.

Patients are usually asked to return at about 6 weeks. If the eye is quiet and the CME has resolved, then the steroid is tapered over the following 2 to 3 months; the patient returns after the taper ends. If this is the first

time a patient has performed a steroid taper, then I wait 4 weeks to bring them back. If they had a previous, more rapid taper, however, I may ask them to return sooner.

THREE SCENARIOS

After another 4 to 6 weeks, the patient is reassessed for recurrent inflammation. At that time, one of three scenarios is encountered.

► Scenario No. 1: No inflammation.

Usually, the eye is quiet, in which case there is nothing more to do.

► Scenario No. 2: Recurrent inflammation.

A subset of patients return with recurrent inflammation at or before this follow-up visit. In the presence of recurrent inflammation, either the steroid taper can be extended, which may not be effective, or the dosing frequency of the topical steroid can be increased while authorization from their insurance company is requested to administer an injectable medication such as the dexamethasone intravitreal implant 0.7 mg (Ozurdex, Allergan).

I instruct patients to stop all topical medications on the day they receive a dexamethasone intravitreal implant and ask them to return in 6 to 8 weeks for an IOP check and anterior segment evaluation. I see patients with elevated IOP or glaucoma sooner to monitor their pressure. In some situations, it is appropriate to start IOP-lowering therapy when the dexamethasone intravitreal implant is placed. Often, patients need just one of these implants and respond well.

► CATARACT SURGERY

► **Scenario No. 3: Rebound inflammation.**

An even smaller subset of patients develops rebound inflammation and may experience a second or third recurrence. In this situation, I may propose a longer-acting intraocular implant such as the fluocinolone acetonide intravitreal implant 0.18 mg (Yutiq, EyePoint Pharmaceuticals).

Patients who have been on long-term topical and/or intraocular therapy often prefer the option of a single injection that lasts 3 years. I do not have any patients who are a full 3 years out from treatment yet, but none of them has experienced a recurrence so far.

Recently, I saw a patient with persistent postoperative inflammation and placed a dexamethasone intravitreal implant after three unsuccessful attempts at tapering topical corticosteroids (a standard

postoperative taper followed by an extended taper over 2 months and then an even longer taper because the patient initially wanted to avoid intravitreal injection). He responded well to the implant but experienced a recurrence of inflammation at the 4-month mark. A second dexamethasone intravitreal implant was placed, followed by the placement of a fluocinolone acetonide intravitreal implant 8 weeks thereafter. Nine months later, he is doing well and has not experienced recurrent inflammation.

CONCLUSION

Given the potential complexity of chronic inflammation after cataract surgery, patients often require targeted evaluation and therapy. When making a referral, it is helpful for the anterior segment surgeon to convey as much information as possible about the case, including preoperative and surgical

details, symptoms, and the results of any tests and imaging. The possibility of endophthalmitis must be considered, but chronic postoperative inflammation is usually idiopathic. In most cases, there is time for the anterior segment surgeon and retina specialist to work collaboratively and ensure that the best decisions are being made for each patient. ■

1. Elliott D, Kim I, Yonekawa Y. Pseudophakic cystoid macular edema (Irvine-Gass syndrome). EyeWiki. Updated May 31, 2022. Accessed August 11, 2022. [https://eyewiki.aao.org/Pseudophakic_Cystoid_Macular_Edema_\(Irvine-Gass_Syndrome\)](https://eyewiki.aao.org/Pseudophakic_Cystoid_Macular_Edema_(Irvine-Gass_Syndrome))
2. Grzybowski A, Sikorski BL, Ascaso FJ, Huerva V. Pseudophakic cystoid macular edema: update 2016. *Clin Interv Aging*. 2016;11:1221-1229.
3. Taravati P, Lam DL, Leveque T, Van Gelder RN. Postcataract surgical inflammation. *Curr Opin Ophthalmol*. 2012;23(1):12-18.

VAIDEHI S. DEDANIA, MD

- Adult and pediatric vitreoretinal surgeon; Associate Professor of Ophthalmology; Director, Vitreoretinal Surgery Fellowship; NYU Langone Health, New York
- vaidehi.dedania@nyulangone.org
- Financial disclosure: Advisor (Allergan, Apellis, Genentech, Regeneron); Consultant (Character Bio)