

## QUALITY OVER QUANTITY: GETTING THE DATA RIGHT

#### **Objective Versus Subjective Data**

The potential for leveraging data in ophthalmology is immense, yet the foundation must be data quality; as the saying goes, "garbage in, garbage out." Ophthalmology benefits from a wealth of objective measurements, distinguishing it from many other medical fields. Subjective metrics, however, such as manifest refraction and defocus curves, must be approached with caution.

#### **Creating and Refining Nomograms**

One of the most straightforward ways to use refractive surgery data is to develop nomograms for laser vision correction. Although third-party companies offer this service, creating nomograms internally can deepen surgeons' understanding of the nuances and noise in their outcomes. Essentially, nomograms illustrate the line of best fit in an attempted-versus-achieved scatterplot; the dispersion of these data points reflects the surgical noise to minimize, thereby improving the R-squared value, which quantifies how well the model's predictions align with observed outcomes of the regression line.

How can this noise be reduced and outcomes be enhanced? The answer is standardization.

# STANDARDIZATION AS A KEY TO CONSISTENCY IN SURGERY

### Lessons From Industry for Standardization

Industrial processes rely on precise sensors and metrics. Components are produced under consistent environmental controls, achieving uniformity and reproducibility. Refractive and cataract surgery can adopt similar standardization practices by drawing on industrial processes and using tools such as the excimer laser.

When applying these principles to LASIK, each surgery should be performed under standardized air conditions (temperature and humidity)

and with consistent surgical techniques, tissue hydration levels, and procedural speed. Observing the impact of these details on cases, including the R-squared value in nomograms, can illustrate how variations in technique affect outcomes.

This attention to consistency—modeled in part after industrial standards—proves essential in high-precision fields such as refractive surgery. As Henry Ford wisely remarked, "If you think of standardization as the best that you know today, but which is to be improved tomorrow, you get somewhere."

### Advancing Cataract Surgery Through Data and Standardization

Robotic cataract surgery is under development, and debate about its potential advantages over manual procedures is ongoing. Femtosecond laser—assisted cataract surgery already provides enhanced reproducibility, precision, and customization for critical steps in cataract surgery.

Increasing the standardization of these procedures, especially when refining IOL constants and formulas, is essential for optimizing the performance of advanced technology IOLs, such as multifocal and extended depth of focus lenses, to deliver greater spectacle independence.

### INTEGRATING CONNECTIVITY AND DATA PLATFORMS

Fully leveraging today's data-rich environment demands connectivity. One need only imagine transforming all paper records and PDF reports from diagnostic devices into a continuous data stream stored in the cloud. In this model, data from each diagnostic station would populate a central, real-time spreadsheet, with thousands of metrics gathered throughout the preoperative workflow.

This comprehensive dataset enables algorithms to guide surgeons in selecting the most appropriate treatment for each case, thereby unlocking the full potential of data-driven, personalized cataract surgery.

Just as oil must be refined, careful curation of data is required to extract their value. Applying advanced algorithms or even machine learning tools can transform raw data into actionable surgical insights—a powerful evolution in personalized cataract surgery.

#### **Overcoming Data Integration Challenges**

Data extraction from ophthalmic equipment presents major challenges because different devices export information using various protocols (eg, Digital Imaging and





Figure 1. A customized dashboard highlighting the most crucial parameters to enhance surgical decision-making.






Figure 2. Measures of image quality for point objects are based on the contrast and compactness of the image.

Communications in Medicine [DICOM], Excel [Microsoft], Access [Microsoft]) and unified nomenclature across brands is often lacking.

Despite these obstacles, it is possible to develop a data platform customized to the equipment within a practice, allowing all relevant parameters to be collected simultaneously. Running a surgical algorithm on these data points could provide surgeons with a streamlined dashboard displaying only the most critical parameters (Figure 1). This could facilitate real-time decision-making and potentially improve patients' visual outcomes.

#### **Developing Standardized Outcome Metrics**

The next step is to standardize metrics for outcome measures such as patient satisfaction and improvements in visual acuity and quality of life. These metrics, however, are often subjective and highly dependent on effective communication between the physician and patient. The ideal scenario is one of "underselling and overdelivering." In essence, patient satisfaction is maximized when their outcomes exceed their expectations.

Interestingly, studies have indicated that patient satisfaction rates following multifocal lens implantation have remained at approximately 90% for the past 3 decades and across multiple generations of lenses and technologies. This stability suggests a high level of proficiency in promoting lens technology but also emphasizes the need for more objective parameters to assess visual quality accurately. 1,2

#### LEVERAGING TECHNOLOGY FOR OBJECTIVE VISUAL QUALITY ASSESSMENT

Fortunately, technological advances now allow objective measurements of visual quality using modern diagnostic tools.

#### **Key Metrics for Image Quality Assessment**

In an ideal optical system, a point object is viewed as a compact, high-contrast retinal image (Figure 2). This representation, known as the point spread function, is calculated as the squared magnitude of the inverse Fourier transform of the pupil function.




Figure 3. Objective defocus curve computed by the Osiris.

For more complex objects, such as gratings, the modulation transfer function quantifies how image contrast varies with spatial frequency, and the phase transfer function describes how the image phase shifts with spatial frequency. Together, the modulation transfer function and phase transfer function compose the optical transfer function of the eye, which is computed as the Fourier transform of the point spread function.

#### **Objective Defocus Curves and Aberrometry Devices**

Collecting these data during follow-up examinations after surgery provides an objective and efficient means of assessing patients' visual quality. These data can serve as crucial input for a machine-learning platform, where linear and nonlinear correlations with preoperative parameters can inform future surgical planning.

Additionally, modern aberrometry devices such as the Osiris (CSO; Figure 3) can generate objective defocus curves through high-quality sensors. These robust feedback data, when integrated into the platform, enhance its learning capacity and may facilitate optimal IOL selection for each eye, based on preoperative factors such as pupil aperture and dynamics.

#### MACHINE LEARNING AND DATA-DRIVEN SURGICAL DECISIONS

After decades of developing high-quality sensors and computing power, the focus of ophthalmology may soon shift to software and information technology. Together with machine learning, these tools offer a path to continuous improvement in the quality of ophthalmic practice by leveraging objective outcomes data to drive strategic advances.

1. Lwowski C, Pawlowicz K, Petermann K, Hemkeppler E, Hinzelmann L, Böhm M, Kohnen T. Visual and patient-reported factors leading to satisfaction after implantation of diffractive extended depth-of-focus and trifocal intraocular lenses I Cotornet Refract Sura 2022-48(4):421-428

2. Chiam PJ, Chan JH, Haider SI, Karia N, Kasaby H, Aggarwal RK. Functional vision with bilateral ReZoom and ReSTOR intraocular lenses 6 months after cataract surgery. J Cataract Refract Surg. 2007;33(12):2057-2061.

#### ANDREA RUSSO, MD, PHD

- **Founder and Medical Director, Gruppo Refrattivo Italiano, Italy**
- andrea.russo@grupporefrattivoitaliano.it
- = Financial disclosure: None